

Experiments with Schemes for Exponential Decay

Hans Petter Langtangen^{1,2} (hpl@simula.no)

¹Center for Biomedical Computing, Simula Research Laboratory

²Department of Informatics, University of Oslo.

Jul 2, 2016

Abstract

This report investigates the accuracy of three finite difference schemes for the ordinary differential equation $u' = -au$ with the aid of numerical experiments. Numerical artifacts are in particular demonstrated.

Contents

1	Mathematical problem	2
2	Numerical solution method	2

3	Implementation	3
4	Numerical experiments	4
5	Error vs Δt	7
	References	9

1 Mathematical problem

We address the initial-value problem

$$u'(t) = -au(t), \quad t \in (0, T], \quad (1)$$

$$u(0) = I, \quad (2)$$

where a , I , and T are prescribed parameters, and $u(t)$ is the unknown function to be estimated. This mathematical model is relevant for physical phenomena featuring exponential decay in time, e.g., vertical pressure variation in the atmosphere, cooling of an object, and radioactive decay.

2 Numerical solution method

We introduce a mesh in time with points $0 = t_0 < t_1 \cdots < t_{N_t} = T$. For simplicity, we assume constant spacing Δt between the mesh points: $\Delta t = t_n - t_{n-1}$, $n = 1, \dots, N_t$.

Let u^n be the numerical approximation to the exact solution at t_n .

The θ -rule [1] is used to solve (1) numerically:

$$u^{n+1} = \frac{1 - (1 - \theta)a\Delta t}{1 + \theta a\Delta t} u^n,$$

for $n = 0, 1, \dots, N_t - 1$. This scheme corresponds to

- The **Forward Euler** scheme when $\theta = 0$
- The **Backward Euler** scheme when $\theta = 1$
- The **Crank-Nicolson** scheme when $\theta = 1/2$

3 Implementation

The numerical method is implemented in a Python function [2] `solver` (found in the `model.py` Python module file):

```
def solver(I, a, T, dt, theta):  
    """Solve  $u' = -a*u$ ,  $u(0) = I$ , for  $t$  in  $(0, T]$  with steps of  
    dt = float(dt) # avoid integer division  
    Nt = int(round(T/dt)) # no of time intervals  
    T = Nt*dt # adjust T to fit time step d  
    u = zeros(Nt+1) # array of  $u[n]$  values  
    t = linspace(0, T, Nt+1) # time mesh  
  
    u[0] = I # assign initial condition  
    for n in range(0, Nt): #  $n=0, 1, \dots, Nt-1$   
        u[n+1] = (1 - (1-theta)*a*dt)/(1 + theta*dt*a)*u[n]  
    return u, t
```

4 Numerical experiments

A set of numerical experiments has been carried out, where I , a , and T are fixed, while Δt and θ are varied. In particular, $I = 1$, $a = 2$, $\Delta t = 1.25, 0.75, 0.5, 0.1$. Figure 1 contains four plots, corresponding to four decreasing Δt values. The red dashed line represent the numerical solution computed by the Backward Euler scheme, while the blue line is the exact solution. The corresponding results for the Crank-Nicolson and Forward Euler methods appear in Figures 2 and 3.




Figure 1: The Backward Euler method for decreasing time step values.




Figure 2: The Crank-Nicolson method for decreasing time step values.




Figure 3: The Forward Euler method for decreasing time step values.

5 Error vs Δt

How the error

$$E^n = \left(\int_0^T (Ie^{-at} - u^n)^2 dt \right)^{\frac{1}{2}}$$

varies with Δt for the three numerical methods is shown in Figure 4.

Observe:

The data points for the three largest Δt values in the Forward Euler method are not relevant as the solution behaves non-physically.

The E numbers corresponding to Figure 4 are given in the table below.

Δt	$\theta = 0$	$\theta = 0.5$	$\theta = 1$
1.25	7.4630	0.2161	0.2440
0.75	0.6632	0.0744	0.1875
0.50	0.2797	0.0315	0.1397
0.10	0.0377	0.0012	0.0335

Summary.

1. $\theta = 1$: $E \sim \Delta t$ (first-order convergence).




Figure 4: Variation of the error with the time step.

2. $\theta = 0.5$: $E \sim \Delta t^2$ (second-order convergence).
3. $\theta = 1$ is always stable and gives qualitatively correct results.
4. $\theta = 0.5$ never blows up, but may give oscillating solutions if Δt is not sufficiently small.
5. $\theta = 0$ suffers from fast-growing solution if Δt is not small enough, but even below this limit one can have oscillating solutions (unless Δt is sufficiently small).

References

- [1] A. Iserles. *A First Course in the Numerical Analysis of Differential Equations*. Cambridge University Press, second edition, 2009.
- [2] H. P. Langtangen. *A Primer on Scientific Programming With Python*. Springer, fifth edition, 2016.